## An Interdisciplinary Approach for Developing Sustainable Grazing Management: Hydrology and Water Quality.

(A05-endale318834-poster)

## Authors:

- D.M. Endale\* USDA-ARS
- D.H. Franklin USDA-ARS
- A.J. Franzluebbers *USDA-ARS*
- J.A. Stuedemann USDA-ARS

## Abstract:

Tall fescue, an important cool-season perennial forage in the humid USA, is usually infected with a fungal endophyte which produces alkaloids toxic to grazing animals. The endophyte/fescue mutualistic association, however, has broader ecological implications. We are evaluating animal, plant, soil, hydrologic and water quality responses of grazing cattle on tall fescue. An endophyte-free and two endophyte-infected fescue types (high or low alkaloid) with broiler litter (BL) or NH4NO3 (AN) fertilization were replicated on two 1-ha paddocks each. Two hayed control paddocks were planted to low-alkaloid fescue and fertilized with NH4NO3. In the first year, nutrient concentrations (NH4-N, NO3-N and PO4-P) were higher from grazed paddocks about 80% of the time. Mean concentrations (skewed by few high values) in mg/L for BL, AN and control treatments were, respectively: NH4-N - 1.2, 0.6, 0.5; NO3-N - 0.3, 0.8, 1.5; PO4-P - 2.0, 0.7, 0.2. Clear fescue treatment effects have not yet emerged. But the high-alkaloid and endophyte-free types under BL had higher offsite nutrient delivery about 70% of the time. The endophyte-free fescue had higher PO4-P concentrations under AN treatment. Treatment effect on runoff is not yet established. In some paddocks 25 to 40% of the rainfall was partitioned into runoff in some events. Data were not complete due to sensor problems.

**Speaker Information**: Dinku Endale, USDA-ARS, 1420 Experiment Station Road, Watkinsville, GA 30677; Phone: 706-769-5631; E-mail: dendale@arches.uga.edu

**Session Information**: Tuesday, November 4, 2003, 4:00 PM-6:00 PM **Presentation Start**: 4:00 PM (Poster Board Number: 204)

**Keywords**: Water quality; Grazing management; Offsite impact; Watershed hydrology